

Enamelast™ Fluoride Varnish 5% Sodium Fluoride Varnish	EN	Enamelast™ Fluoride Varnish 5%iger Natriumfluoridlack	DE	Enamelast™ Fluoride Varnish Vernis à 5 % de fluorure de sodium	FR	Enamelast™ Fluoride Varnish 5% Sodium Fluoride Vernis	NL	Enamelast™ Fluoride Varnish Vernice al fluoruro al 5%	IT	Enamelast™ Fluoride Varnish Barniz de fluoruro de sodio al 5 %	ES	Enamelast™ Fluoride Varnish Verniz de fluoruro de sódio a 5 %	PT	Enamelast™ Fluoride Varnish 5 % natriumfluoridlakk	SV
Product Description															
Enamelast™ fluoride varnish is a flavored, xylitol-sweetened, 5% sodium fluoride in a resin carrier delivered in a 1.2 ml syringe or a 0.4 ml unit dose blister pack.															
For all products described, carefully read and understand all instructions and SDS information prior to use.															
Delivery Form(s)															
<ul style="list-style-type: none"> • 1.2 ml delivery syringe ◦ SoftEZ™ tip • 0.4 ml unit dose pack ◦ Application brush 															
Indications for Use/Intended Purpose															
Enamelast™ fluoride varnish is 5% sodium fluoride in a varnish carrier which produces a mechanical occlusion of the dentinal tubules in the treatment of tooth hypersensitivity.															
Contraindications															
<ul style="list-style-type: none"> • Do not use on patients with ulcerative gingivitis or stomatitis. • Do not use on patients with a known sensitivity to ingredients including clophony (klofonium) and rosan. If allergic reaction, dermatitis, or rash develops, consult a physician. • For patients or users with allergy concerns, refer to product allergen document available at www.ultradent.com. If an allergic reaction is observed, rinse exposed area thoroughly with water and have the patient consult their physician. 															
Warnings and Precautions															
<ul style="list-style-type: none"> • Use only recommended tips. • Tips are single use only. • Patient, clinician, and assistant must wear protective eyewear. • In rare instances, edematous swelling has been reported after extensive fluoride varnish treatment and dyspnea has been reported in asthmatic patients. If this occurs, thoroughly brush teeth to remove the varnish film and have patient contact their medical provider. • Some patients may experience nausea after application. If nausea persists, thoroughly brush to remove the varnish film. • Do not use more than half of the 1.2 ml syringe of fluoride varnish per patient per application. • Do not use with other topical fluoride treatments the same day and instruct patient to refrain from using oral fluoride supplements for 4 (four) days after treatment. • Do not remove cap from the syringe or foil lid from the unit dose until ready to use. 															
Stepwise Instructions															
Preparation															
1) Thoroughly brush teeth, rubber cup polish, or wipe teeth with gauze prior to placement.															
2) For syringe:															
<ul style="list-style-type: none"> a. Securely attach recommended delivery tip and verify the flow, color and consistency of the product before applying by expressing a small amount onto a mixing pad or gauze. Discard any clear liquid that is expressed. If resistance is met, replace tip and recheck. b. Open blister pack by peeling back and removing the foil lid. Using the disposable application brush, lightly stir the varnish. 															
3) For unit dose:															
4) Open blister pack by peeling back and removing the foil lid. Using the disposable application brush, lightly stir the varnish.															
Gebrauch															
1) Zu behandelnder trockener Bereich.															
2) Setzen Sie eine dünne, gläzende Schicht Enamelast Fluoridlack auf viele Oberflächen des Zähns mit dem Einwegspatel.															
3) Auf Spülung warten.															
4) Auf Spülung warten.															
Verarbeitung															
1) Putzen Sie die Zähne gründlich, polieren Sie die Gummikappe oder wischen Sie die Zähne mit dem Einweg mit Gaze ab.															
2) Auf Spülung warten.															
3) Setzen Sie die empfohlene Abgabespitze sicher an und überprüfen Sie den Fluss, die Farbe und die Konsistenz des Produkts, bevor Sie es anwenden. Siehe eine kleine Menge auf einer Armchlicko oder einer Gaze ausdrücken. Ausgedrückte klare Flüssigkeit entspricht. Bei Widerstand die Kanüle entfernen und neu einsetzen.															
4) Für Einheitsdosier:															
a. Öffnen Sie die Blisterpackung, indem Sie den Foliendeckel abziehen und entfernen. Mit dem Einweg-Applikatorspatel die Löffel abziehen.															
b. Auf Spülung warten.															
c. Auf Spülung warten.															
d. Auf Spülung warten.															
e. Auf Spülung warten.															
f. Auf Spülung warten.															
g. Auf Spülung warten.															
h. Auf Spülung warten.															
i. Auf Spülung warten.															
j. Auf Spülung warten.															
k. Auf Spülung warten.															
l. Auf Spülung warten.															
m. Auf Spülung warten.															
n. Auf Spülung warten.															
o. Auf Spülung warten.															
p. Auf Spülung warten.															

